

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

In UNITED STATES PATENT AND TRADE MARK OFFICE

PROVISION PATENT APPLICATION

ENTITLED:

Stable Gabapentin having pH within a Controlled Range

INVENTORS:

Claude Singer  
Gideon Pilarsky  
Michael Pesachovitch

Attorney Docket No. 1662/52802

## Stable Gabapentin having pH within a Controlled Range

### Cross-reference to Related Applications

This invention relates to PCT Application No. WO  
5 98/28255, filed July 2, 1998, also assigned to the  
assignee of the present invention and incorporated  
herein by reference; the present invention also claims  
priority to U.S. Provisional Application No. 60/211,966,  
filed June 16, 2000.


10

### Field of the Invention

The present invention relates to a pharmaceutical  
composition containing therapeutically effective amount  
of gabapentin and its derivatives in combination with  
15 effective carriers. More particularly, the present  
invention relates to a stable composition and a process  
for manufacturing pure and stable gabapentin having a pH  
in the range of 6.8 to 7.3.

### Background of the Invention

20 Gabapentin is 1-(aminomethyl)-1-cyclohexanecarboxylic  
acid, having the chemical structure of formula I:



Gabapentin is used for treating cerebral diseases  
such as epilepsy, faintness attacks, hypokinesia and  
25 cranial traumas. United States Patent No. 4,024,175 to  
Satzinger et al., incorporated herein by reference,  
discloses that gabapentin of formula (I) shows

U.S. Patent Application


Filed: June 15, 2001

Inventors: Claude Singer, Gideon Pilarsky and Michael Pesachovitch

Attorney Docket No. 1662/52802

hypothermal and, in some cases, narcosis-potentiating or sedating properties as well as protective effect against cardiozole cramp in animals. Finally, gabapentin has been found especially useful in treating geriatric  
5 patients. As such, there has been a need for producing pure and stable gabapentin.

United States Patent No. 6,054,482 to Augart et al. discloses that preparation and long-term storage of gabapentin presents several problems since (i) during  
10 the preparation the compounds shows considerable variations without apparent reason, and (ii) the long-term storage of even very pure gabapentin showed differing stabilities with progressively long storage times. Augart further discloses that the toxic lactam  
15 compound of formula (II)



forms during the preparation and storage of gabapentin. According to Augart, because the lactam has  
20 higher toxicity than gabapentin, its presence in gabapentin should be limited if not eliminated. To combat the lactam formation and provide product stability, Augart stresses the importance of (i) starting with gabapentin raw material that contains 0.5%  
25 or less of corresponding lactam, (ii) not allowing the anion of a mineral acid in the composition to exceed 20 ppm, and (iii) using a specifically selected adjuvant

U.S. Patent Application  
Filed: June 15, 2001  
Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch  
Attorney Docket No. 1662/52802

that is not adverse to gabapentin stability.

According to Augart, the following adjuvants (or excipients) had no noticeable influence on the stability of gabapentin, and as such, they were taught to be acceptable adjuvants for use with gabapentin: hydroxypropylmethylcellulose, polyvinylpyrrolidone, crospovidon, poloxamer 407, poloxamer 188, sodium starch glycolate, copolyvidone, maize starch, cyclodextrin, lactose, talc, as well as co-polymers of dimethylamino-methacrylic acid and neutral methacrylic acid ester.

Conversely, Augart discloses that the following adjuvants reduce the stability of gabapentin and should be avoided: modified maize starch, sodium croscarmellose, glycerol behenic acid ester, methacrylic acid co-polymers (types A and C), anion exchangers titanium dioxide and silica gels such as Aerosil 200.

The composition and method disclosed in Augart are industrially impractical and technically unnecessary. It has now been found that Augart's reliance on maintaining the anion of a mineral acid as not exceeding 20 ppm is misplaced. Thus, gabapentin and pharmaceutical formulations of gabapentin can be prepared and stored such that initially they do not contain more than 0.5% of the lactam and even after one year of storage at 25 °C and 60% atmospheric humidity, the conversion of gabapentin to its corresponding lactam does not exceed 0.2% by weight of gabapentin. That is, gabapentin and pharmaceutical formulations of gabapentin have been found to be stable even though such formulations do not meet Augart's requirements (ii) and (iii).

U.S. Patent Application  
Filed: June 15, 2001  
Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch  
Attorney Docket No. 1662/52802

The specific mineral acid disclosed by Augert is hydrochloric acid (column 3, lines 61-63; column 5, lines 24-29; examples 1 and 2). The specification states, in particular

5           The active materials of formula (I) [including gabapentin] must be prepared as highly purified, nonderivatized free amino acids, for example, from the corresponding hydrochloride by ion exchange.  
10          The proportion of remaining hydrochloride admixtures should thereby not exceed 20 ppm.

(Column 5, lines 24-29).

20 ppm of gabapentin hydrochloride corresponds to roughly 3 ppm of chloride ion, due to the higher  
15          molecular weight of gabapentin.

Augert's claims require gabapentin with "less than 20 ppm of the anion of a mineral acid", e.g. chloride.

#### Summary of the Invention

Accordingly, the present invention relates to a  
20          pharmaceutical composition containing a pharmaceutically effective amount of gabapentin having a pH in the range of 6.8 to 7.3 and which initially contains less than 0.5% of a corresponding lactam and after one year of storage at 25 °C and 60% atmospheric humidity the  
25          conversion of gabapentin to its corresponding lactam does not exceed 0.2% by weight of gabapentin.

The present invention also relates to a process for preparing a stable pharmaceutical formulation containing gabapentin having pH in the range of 6.8-7.3, more  
30          preferably in the range of 7.0-7.2, initially containing

U.S. Patent Application

Filed: June 15, 2001

Inventors: Claude Singer, Gideon Pilarsky and Michael Pesachovitch

Attorney Docket No. 1662/52802

less than 0.5% of a corresponding lactam and after storage for one year at 25 °C and 60% atmospheric humidity the conversion of gabapentin to its corresponding lactam does not exceed 0.2% by weight of  
5 gabapentin.

Detailed Description of the Preferred Embodiments

The subject invention will now be described in greater detail for preferred embodiments of the invention, it being understood that these embodiments  
10 are intended only as illustrative examples and the invention is not to be limited thereto.

As will be illustrated through exemplary embodiments 1-16, gabapentin may be prepared from the hydrochloride salt of gabapentin (gabapentin  
15 hydrochloride) and that in purified form gabapentin may have a pH in the range of 6.8-7.3, and preferably in the range of 7.0-7.2. The gabapentin formulation may also contain more than 20 ppm of chloride ion in the composition as measured by the amount of chloride ion in  
20 the composition.

Exemplary embodiments 17-19 illustrate formulations of gabapentin containing varying amounts of chloride ion, some of which are greater than 20 ppm and some less, and all of which initially contain less than 0.5%  
25 of lactam and after one year of storage at 25 °C and 60% humidity, the conversion of gabapentin to its corresponding lactam is measured not to exceed 0.2% by weight of gabapentin.

Commonly known adjuvants (also referred to as  
30 excipients) which can be utilized in a gabapentin

U.S. Patent Application

Filed: June 15, 2001

Inventors: Claude Singer, Gideon Pilarsky and Michael Pesachovitch

Attorney Docket No. 1662/52802

formulation of the present invention may include for example, ~~modified maize starch, sodium croscarmellose, titanium dioxide,~~ and (silica gels such as Aerosil 200. Hydroxypropylmethylcellulose, polyvinylpyrrolidone,

5 crosppovidon, poloxamer 407, poloxamer 188, sodium starch glycolate, copolyvidone, maize starch, cyclodexterin, lactose, talc, co-polymers of dimethylamino-methacrylic acid and neutral methacrylic acid ester may also be used. The list of adjuvants is not an exhaustive list.  
10 and it would be within the scope of the claimed invention to use any known adjuvant that would behave similar to those enumerated herein.

Certain specific representative embodiments of the invention are described in detail below, the materials,  
15 apparatus and process steps being understood as examples that are intended for illustrative purposes only. Consequently, it will be noted that the invention is not intended to be limited to the methods, materials, conditions, precess parameters, apparatus and the like  
20 specifically recited herein.

In the examples below chloride ion concentration is measured by any commonly known method, such as for example, by titration with  $\text{AgNO}_3$ , pH electrode or chromatography.

25 EXAMPLE 1

The following raw material were used:

Gabapentin hydrochloride	18.2 g
Isopropanol for dissolution	160 ml
Active carbon SX1	1.1 g
30 Ethylacetate	268 ml



U.S. Patent Application  
Filed: June 15, 2001  
Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch  
Attorney Docket No. 1662/52802

Tributylamine	19.5 g
Methanol for washing	23 ml

A) Preparation of Crude gabapentin

5 Gabapentin hydrochloride was dissolved in 130 ml of dry isopropanol at 25°C by mixing. Next, 1.1 grams of active carbon was added and the suspension was heated to 40°C and maintained at this temperature for 2 hours. The suspension was then filtered at 40°C and the filter  
10 cake was washed twice with additional 15 ml of isopropanol each time. The washings were added to the already separated solution of gabapentin hydrochloride in isopropanol. The solution was concentrated to dryness in vacuum (Approximately 10 mm Hg) to a constant  
15 weight. The temperature of the heating bath was maintained (maximally) at 35°C during this operation. Thereafter, 245 ml of ethylacetate was added to the dry residue of gabapentin hydrochloride and the solution was mixed. After half an hour of mixing at 25°C, an amount  
20 of 19.5 grams of tributylamine was added during the subsequent 30 minutes. The mixing continued for an additional two hours at the same temperature.

The gabapentin base which was formed during this operation was separated from the suspension through  
25 filtration. The filter cake was washed with 23 ml of ethylacetate and 23 ml of methanol to give crude gabapentin.

B) Gabapentin Purification

The following raw material were used:

30	Methanol for suspending	52.5 ml
----	-------------------------	---------

U.S. Patent Application  
Filed: June 15, 2001  
Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch  
Attorney Docket No. 1662/52802

Methanol for washing 2x15 ml

Wet crude gabapentin prepared according to Step A was suspended in 52.5 ml of methanol for 14 hours at approximately 25°C and stirred. Thereafter, the solid  
5 gabapentin was separated from the suspension by filtration. The filter cake was washed twice with 15 ml of methanol and than dried under vacuum giving pure gabapentin. The yield was 72%.

The following data regarding the chlorine anion  
10 content of the above-prepared gabapentin were obtained:

**TABLE 1 - Anion content and pH values  
after the reslurry in methanol**

Run	Cl <sup>-</sup> (ppm)	pH
A	4	6.94
B	20	7.01
C	< 5	7.04
D	40	6.97
E	35	6.92
F	15	6.84

15

20

Gabapentin purified according to these procedures contains less than 0.5% lactam as measured by HPLC vs. standard. After a year of storage at 25°C and 60% relative humidity, the conversion of gabapentin to its  
25 corresponding lactam is measured not to exceed 0.2% by weight of gabapentin.

For a better control of the pH of pure gabapentin several basic agents were added. Some examples of added basic agents are given in the following Examples.

30 **EXAMPLE 2**

The following raw material were used:

U.S. Patent Application  
Filed: June 15, 2001  
Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch  
Attorney Docket No. 1662/52802

Methanol for suspending	52.5 ml
Methanol for washings	2x15 ml
Tributylamine	~0.3 equivalents

The wet crude gabapentin(as in Step 1A) was  
5 suspended in 52.5 ml of methanol for 14 hours and at  
25°C and stirred. Tributylamine was added to the  
suspension. After 14 hours of stirring the solid  
gabapentin was separated from the suspension by  
filtration. The filter cake was then washed twice, each  
10 time with 15 ml of methanol and than dried under vacuum  
resulting in pure gabapentin with a yield of 87%, pH of  
7.15 and chlorine anion content of 50 ppm. Gabapentin  
so prepared initially contained less than 0.5% by weight  
of lactam, and, after a year of storage at 25°C and 60%  
15 relative humidity, the conversion of gabapentin to its  
corresponding lactam is measured not to exceed 0.2% by  
weight of gabapentin.

### EXAMPLE 3

The following raw material were used:

20	Methanol for suspending	52.5 ml
	Methanol for washing	2x15 ml
	Sodium methoxide	~0.001 equivalents

The wet crude gabapentin (as in Example 1, step A)  
was suspended in 52.5 ml of methanol for 14 hours and  
25 kept at 25°C. Sodium methoxide was added to the  
suspension. After 14 hours of stirring, the solid  
gabapentin was separated from the suspension by  
filtration. The filter cake was then washed twice with

U.S. Patent Application  
Filed: June 15, 2001  
Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch  
Attorney Docket No. 1662/52802

15 ml of methanol, then dried under vacuum, resulting in  
pure gabapentin having a yield of 85%, pH of 6.8, and  
chlorine anion content of 50 ppm. Gabapentin so prepared  
contained less than 0.5% by weight of lactam, and, after  
5 a year of storage at 25°C and 60% relative humidity, the  
conversion of gabapentin to its corresponding lactam is  
measured not to exceed 0.2% by weight of gabapentin.

It should be noted that the solvents and the base  
used in Example 1A were not unique. In addition, it  
10 should be noted that in Examples 4-9 gabapentin pure was  
always prepared as in Example 1B and the results (Cl<sup>-</sup>  
content and yield) refer to gabapentin pure.

#### EXAMPLE 4

The following raw material were used:

15	Gabapentin hydrochloride (100%)	18.2 g
	Isopropanol for dissolution	160 ml
	Active carbon SX1	1.1 g
	Tributylamine	19.5 g
	Methanol for washing	23 ml

20 In this Example, gabapentin hydrochloride was  
dissolved in 130 ml of dry isopropanol at 25°C. Then  
1.1 grams of active carbon was added and the suspension  
was heated to 40° C and maintained at this temperature  
for 2 hours. The suspension was filtered at 40° C and  
25 the filter cake was then washed twice, each time with an  
additional 15 ml of isopropanol. The washings were  
added to the already separated solution of gabapentin  
hydrochloride in isopropanol. After half an hour of

U.S. Patent Application

Filed: June 15, 2001

Inventors: Claude Singer, Gideon Pilarsky and Michael Pesachovitch

Attorney Docket No. 1662/52802

mixing at 25° C, 19.5 grams of tributylamine was added during half an hour and the mixing was continued for two hours at the same temperature. The formed gabapentin base was separated from the suspension by filtration and  
5 washed with 23 ml of methanol to give gabapentin crude. After reslurry as in Example 1B gabapentin pure was obtained at a yield of 58.8% and chloride anion content of 7 ppm Cl<sup>-</sup>.

Gabapentin so prepared contained less than 0.5% by  
10 weight of lactam, and, after a year of storage at 25°C and 60% relative humidity, the conversion of gabapentin to its corresponding lactam is found not to exceed 0.2% by weight of gabapentin.

EXAMPLE 5

15 The following raw material were used:

Gabapentin hydrochloride (100%)	18.2 g
Isopropanol for dissolution	160 ml
Active carbon SX1	1.1 g
Ethylacetate	268 ml
20 Trihexylamine	28.3 g
Methanol for washing	23 ml

Gabapentin hydrochloride was dissolved in 130 ml dry isopropanol at 25° C by mixing, then 1.1 g of active carbon was added and the suspension was heated to 40° C  
25 and maintained for two hours at 40° C. The suspension was filtered at 40°C and the filter cake was washed twice with additional 15 ml of isopropanol each time. The washings were added to the already separated

U.S. Patent Application

Filed: June 15, 2001

Inventors: Claude Singer, Gideon Pilarsky and Michael Pesachovitch

Attorney Docket No. 1662/52802

solution of gabapentin hydrochloride in isopropanol.

The solution was concentrated to dryness in vacuum

(approximately 10 mm Hg) to constant weight. The

temperature of the heating bath was maintained at

5 maximum 35° C during this operation. Next, 245 ml of

ethylacetate was added to the dry residue of gabapentin

hydrochloride and the mixing was started. After half an

hour of mixing at 25°C, an amount of 28.3 grams of

trihexylamine was added during half an hour and the

10 mixing was continued for an additional two hours at the

same temperature. The formed gabapentin base was

separated from the suspension by filtration. The filter

cake was washed with 23 ml of ethylacetate and 23 ml of

methanol to give gabapentin crude. After reslurry as in

15 Example 1B, gabapentin pure was obtained having a yield

of 75.0% and chloride anion content of 213 ppm.

#### EXAMPLE 6

The following raw material were used:

Gabapentin hydrochloride (100%)	18.2 g
---------------------------------	--------

20 Isopropanol for dissolution	160 ml
--------------------------------	--------

Active carbon SX1	1.1 g
-------------------	-------

Ethylacetate	268 ml
--------------	--------

Tripropylamine	15 g
----------------	------

Methanol for washing	23 ml
----------------------	-------

25 Gabapentin hydrochloride was dissolved in 130 ml dry isopropanol at 25°C by mixing, then 1.1 g of active carbon was added and the suspension was heated to 40°C and maintained during two hours at 40° C. The

U.S. Patent Application  
Filed: June 15, 2001  
Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch  
Attorney Docket No. 1662/52802

suspension was filtered at 40° C and the filter cake was washed twice with additional 15 ml of isopropanol each time. The washings were added to the already separated solution of gabapentin hydrochloride in isopropanol.

5 The solution was concentrated to dryness in vacuum (~10 mm Hg) to constant weight. The temperature of the heating bath was maintained at maximum 35° C during this operation. Next, 245 ml of ethylacetate was added to the dry residue of gabapentin hydrochloride and mixing  
10 commenced. After half an hour of mixing at 25° C, fifteen grams of tripropylamine was added during half an hour and the mixing was continued for two hours at the same temperature. The formed gabapentin base was separated from the suspension by filtration. The filter  
15 cake was washed with 23 ml of ethylacetate and 23 ml of methanol to give gabapentin crude. After a reslurry process, as in Example 1B, gabapentin pure was obtained having a yield of 68.0% and chloride anion content of 142 ppm.

20 EXAMPLE 7

The following raw material were used:

	Gabapentin hydrochloride (100%)	18.2 g
	Isopropanol for dissolution	160 ml
	Active carbon SX1	1.1 g
25	Acetonitrile	268 ml
	Tributylamine	19.5 g
	Methanol for washing	23 ml

U.S. Patent Application  
Filed: June 15, 2001  
Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch  
Attorney Docket No. 1662/52802

Gabapentin hydrochloride was dissolved in 130 ml of dry isopropanol at 25° C by mixing, then 1.1 g of active carbon was added and the suspension was heated to 40° C and maintained for two hours at 40° C. The suspension  
5 was filtered at 40° C and the filter cake was washed twice with additional 15 ml of isopropanol. The washings were added to the already separated solution of gabapentin hydrochloride in isopropanol. The solution was concentrated to dryness in vacuum (~10 mm Hg) to  
10 constant weight. The temperature of the heating bath was maintained at a maximum temperature of 35°C during this operation. Next, 245 ml of acetonitrile was added to the dry residue of gabapentin hydrochloride and mixing commenced. After half an hour of mixing at 25°  
15 C, an amount of 19.5 g of tributylamine was added during 30 minutes and the mixing was continued for two hours at the same temperature. The formed gabapentin base was separated from the suspension by filtration. The filter cake was washed with 23 ml of acetonitrile and 23 ml of  
20 methanol to give gabapentin crude. After reslurry as in Example 1B, gabapentin pure was obtained having a yield of 67.8%, and anion content of 142 ppm.

#### EXAMPLE 8

The following raw material were used:

25	Gabapentin hydrochloride (100%)	18.2 g
	Isopropanol for dissolution	160 ml
	Active carbon SX1	1.1 g
	Dimethylcarbonate	268 ml
	Tributylamine	19.5 g



U.S. Patent Application  
Filed: June 15, 2001  
Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch  
Attorney Docket No. 1662/52802

Methanol for washing 23 ml

Gabapentin hydrochloride was dissolved in 130 ml of dry isopropanol at 25° C by mixing, then 1.1 g of active carbon was added and the suspension was heated to 40°C and maintained at 40° C for two hours. The suspension was filtered at 40° C and the filter cake was washed twice with additional 15 ml of isopropanol. The washings were added to the already separated solution of gabapentin hydrochloride in isopropanol. The solution was concentrated to dryness in vacuum (~10 mm Hg) to constant weight. The temperature of the heating bath was maintained at maximum of 35° C during this operation. Next, 245 ml of dimethylcarbonate was added to the dry residue of gabapentin hydrochloride and the mixing was started. After half an hour of mixing at 25° C, an amount of 19.5 g of tributylamine was added during half an hour and the mixing was continued for two hours at the same temperature. The formed gabapentin base was separated from the suspension by filtration. The filter cake was washed with 23 ml of dimethylcarbonate and 23 ml of methanol to give gabapentin crude. After reslurry as in Example 1B, gabapentin pure was obtained, having a yield of 57.9%, and anion content of 142 ppm.

#### EXAMPLE 9

The following raw material were used:

Gabapentin hydrochloride (100%)	18.2 g
Isopropanol for dissolution	160 ml
Active carbon SX1	1.1 g
Isopropylacetate	268 ml

U.S. Patent Application

Filed: June 15, 2001

Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch

Attorney Docket No. 1662/52802

Tributylamine 19.5 g

Methanol for washing 23 ml

Gabapentin hydrochloride is dissolved in 130 ml of  
dry isopropanol at 25° C by mixing, then 1.1 g of active  
5 carbon was added and the suspension was heated to 40°C  
and maintained for two hours at 40°C. The suspension  
was filtered at 40°C and the filter cake was washed  
twice with additional 15 ml of isopropanol each time.  
The washings were added to the already separated  
10 solution of gabapentin hydrochloride in isopropanol. The  
solution was concentrated to dryness in vacuum (~10 mm  
Hg) to constant weight. The temperature of the heating  
bath was maintained at maximum 35° C during this  
operation. Next, 245 ml of isopropylacetate was added  
15 to the dry residue of gabapentin hydrochloride and  
mixing commenced. After half an hour of mixing at 25°C,  
an amount of 19.5 g of tributylamine was added during  
half an hour and the mixing was continued for two hours  
at the same temperature. The formed gabapentin base was  
20 separated from the suspension by filtration. The filter  
cake was washed with 23 ml of isopropylacetate and 23 ml  
of methanol to give gabapentin crude. After reslurry as  
in Example 1B, gabapentin pure was obtained having a  
yield of 57.9% and an anion content of 142 ppm.

25 EXAMPLE 10

(The neutralization reaction as in Example 1, however,  
the reslurry in methanol is replaced by a  
crystallization in methanol.)

The following raw material were used:

30 Methanol for dissolution 180 ml

U.S. Patent Application  
Filed: June 15, 2001  
Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch  
Attorney Docket No. 1662/52802

Methanol for washing          2x12 ml

The gabapentin crude (Step 1A) was suspended in 180 ml of methanol at 25° C. The suspension was heated while mixing to 55°C when gabapentin was dissolved. The solution was then cooled slowly for an hour to 25°C. At 25° C the solution was concentrated to a volume of 50 ml. The suspension was stirred for twelve hours at 25°C. After 12 hours, the solid gabapentin was separated from the suspension by filtration. The filter cake was washed twice with 12 ml of methanol then dried under vacuum to give gabapentin pure (yield: 72%). Following Cl<sup>-</sup> contents of gabapentin and pH values were obtained and tabulated in TABLE 2 as follows:

**TABLE 2 - Anion content and pH values for crystallization in methanol**

Run	Cl <sup>-</sup> (ppm)	pH
A	4	6.94
B	< 5	7.2
C	150-200	6.9

For a better control of the pH of gabapentin pure several basic agents were added. Some examples of added basic agents are given in the following examples

**EXAMPLE 11**

The following raw material were used:

Methanol for suspending          52.5 ml

Methanol for washings          2x15 ml

U.S. Patent Application  
Filed: June 15, 2001  
Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch  
Attorney Docket No. 1662/52802

Tributylamine ~0.34 equivalents

The gabapentin crude was suspended in 180 ml of methanol at 25°C. The suspension was then heated, while mixing, to 55°C when gabapentin was dissolved.

5 Tributylamine was added to the solution and the solution was cooled slowly during an hour to a temperature of 25° C. At 25° C the solution was concentrated to a volume of 50 ml. The suspension was stirred for twelve hours at 25° C. After 12 hours the solid gabapentin was  
10 separated from the suspension by filtration. The filter cake was washed twice with 12 ml of methanol and then dried under vacuum to give gabapentin pure having a yield of 81.4%, pH of 7.25 and chlorine anion content of 35 ppm.

15 Gabapentin so prepared initially contained less than 0.5% by weight of lactam, and, after a year of storage at 25°C and 60% relative humidity, the conversion of gabapentin to its corresponding lactam is measured not to exceed 0.2% by weight of gabapentin.

20 EXAMPLE 12

The following material were used:

Methanol for suspending 52.5 ml

Methanol for washings 2x15 ml

Sodium methoxide ~0.001 equivalents

25 Crude gabapentin was suspended in 180 ml of methanol at 25°C. The suspension was heated under mixing to 55° C when gabapentin was dissolved. Sodium methoxide was added to the solution and the solution was cooled slowly during one hour to 25° C. At 25°C the

U.S. Patent Application  
Filed: June 15, 2001  
Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch  
Attorney Docket No. 1662/52802

solution was concentrated to a volume of 50 ml. The  
suspension was stirred for twelve hours at 25°C. After  
12 hours the solid gabapentin was separated from the  
suspension by filtration. The filter cake was washed  
5 twice with 12 ml of methanol, then dried under vacuum to  
give gabapentin pure at a yield of 81.4%, pH of 7.08 and  
anion content of Cl<sup>-</sup> 20 ppm.

Gabapentin so prepared contained less than 0.5% by  
weight of lactam, and, after a year of storage at 55°C  
10 and 50% relative humidity, the amount of lactam remained  
less than 0.5% by weight. After a year of storage at  
25°C and 60% relative humidity, the conversion of  
gabapentin to its corresponding lactam is found not to  
exceed 0.2% by weight of gabapentin.

15 EXAMPLE 13

The following material were used:

Methanol for suspending 52.5 ml

Methanol for washings 2x15 ml

Sodium bicarbonate ~0.05 equivalents

20 Crude gabapentin was suspended in 180 ml of  
methanol at 25° C. The suspension was heated under  
mixing to 55°C when gabapentin was dissolved. Sodium  
bicarbonate was added to the solution and the solution  
was cooled slowly for one hour to 25°C. At 25°C the  
25 solution was concentrated to a volume of 50 ml. The  
suspension was stirred for twelve hours at 25°C. After  
12 hours the solid gabapentin was separated from the  
suspension by filtration. The filter cake was washed  
twice with 12 ml of methanol, then dried under vacuum to

U.S. Patent Application  
Filed: June 15, 2001  
Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch  
Attorney Docket No. 1662/52802

give gabapentin pure having a yield of 72.4%, pH of 7.28  
and anion ( $\text{Cl}^-$ ) content of 20 ppm.

Gabapentin so prepared contained less than 0.5% by  
weight of lactam, and, after a year of storage at 25°C  
5 and 50% relative humidity, the amount of lactam remained  
less than 0.5% by weight. After a year of storage at  
25°C and 60% relative humidity, the conversion of  
gabapentin to its corresponding lactam is found not to  
exceed 0.2% by weight of gabapentin.

10 EXAMPLE 14

The following raw material were used:

Methanol for suspending	52.5 ml
Methanol for washing	2x15 ml
Tetramethylammoniumhydroxide	-0.002 equivalents

15 Crude gabapentin was suspended in 180 ml of  
methanol at 25° C. The suspension was heated under  
mixing to 55° C when gabapentin was dissolved.  
Tetramethylammoniumhydroxide was added to the solution  
and the solution was cooled slowly for one hour to 25°C.  
20 At 25° C the solution was concentrated to a volume of 50  
ml. The suspension was stirred for 12 hours at 25°C.  
After 12 hours the solid gabapentin was separated from  
the suspension by filtration. The filter cake was  
washed twice with 12 ml of methanol than dried under  
25 vacuum to give gabapentin pure having a yield of 75.8%,  
pH of 7.03 and anion content of ( $\text{Cl}^-$ ) 20 ppm.

Gabapentin so prepared initially contained less  
than 0.5% by weight of lactam.

U.S. Patent Application  
Filed: June 15, 2001  
Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch  
Attorney Docket No. 1662/52802

**EXAMPLE 15**

The following raw material were used:

	Methanol for suspending	52.5 ml
	Methanol for washing	2x15 ml
5	Tetrabutylammoniumhydroxide	~0.002 equivalents

Crude gabapentin was suspended in 180 ml of methanol at 25° C. The suspension was heated under mixing to 55°C when gabapentin was dissolved. Tetrabutylammoniumhydroxide was added to the solution and the solution was cooled slowly during one hour to 25°C. At 25°C the solution was concentrated to a volume of 50 ml. The suspension was stirred for 12 hours at 25°C. After 12 hours the solid gabapentin was separated from the suspension by filtration. The filter cake was washed twice with 12 ml of methanol, then dried under vacuum to give gabapentin pure having a yield of 77.6%, pH of 7.22 and anion (Cl<sup>-</sup>) content of 20 ppm.

**EXAMPLE 16**

The following raw material were used:

20	Methanol for suspending	52.5 ml
	Methanol for washing	2x15 ml
	Sodiumtetraborate	~0.05 equivalents

Crude gabapentin was suspended in 180 ml of methanol at 25° C. The suspension was heated under mixing to 55°C when gabapentin was dissolved. Sodiumtetraborate was added to the solution and the solution was cooled slowly for one hour to 25°C. At

U.S. Patent Application

Filed: June 15, 2001

Inventors: Claude Singer, Gideon Pilarsky and Michael Pesachovitch

Attorney Docket No. 1662/52802

25°C the solution was concentrated to a volume of 50ml. The suspension was stirred for 12 hours at 25°C. After 12 hours the solid gabapentin was separated from the suspension by filtration. The filter cake was washed  
5 twice with 12 ml of methanol and then dried under vacuum to give gabapentin pure having a yield of 75%, pH of 7.17 and anion content (Cl<sup>-</sup>) of 10 ppm.

**EXAMPLE 17**

The following gabapentin tablet formulation is  
10 prepared using gabapentin containing chloride ion ranging from 5 to 40 ppm and pH in the range of 6.84 - 7.04 according to Example 1. The following material is used:

Ingredients	Amounts
15 gabapentin	125 g
Corn Starch NF	200 g
Cellulose, Microcrystalline	46 g
Sterotex Powder HM	4 g
20 Purified Water	q.s. or 300 ml

Combine corn starch, cellulose, and gabapentin together in a mixer and mix for 2-4 minutes. Add water to this combination and mix for an addition 1-3 minutes. The resulting mix is spread on trays and dried in  
25 convection oven at 45-55 °C until a moisture level of 1 to 2% is obtained. The dried mix is then milled and added back to the mill mixture and the total is blended for additional 4-5 minutes. Compressed tables of 150 mg, 375 mg and 750 mg are formed using appropriate  
30 punches from the total mix.



U.S. Patent Application  
Filed: June 15, 2001  
Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch  
Attorney Docket No. 1662/52802

The formulation is measured to contain less than 0.5% lactam and after one year of storage at 25 °C and 60% atmospheric humidity, the conversion of gabapentin to its corresponding lactam is measured not to exceed 5 0.2% by weight of gabapentin.

#### EXAMPLE 18

Gabapentin of Example 2 (having chloride ion content of 50 ppm and pH of 7.15) is used to formulate tablets as in EXAMPLE 17, except that corn starch is  
10 replaced in each sample by one of the following adjuvants: pregelatinized starch, croscarmellose sodium, silica gel, titanium dioxide, talc, modified maize starch and maize starch.

The resulting gabapentin tablet of each sample is  
15 initially measured to have 0.5 % by weight of a corresponding lactam, more than 50 ppm of chloride anion, and pH exceeding 6.8. The tablet is stored for one year at 25 °C and 60% atmospheric humidity and the conversion of gabapentin to its corresponding lactam is  
20 found not to exceed 0.2% by weight of gabapentin.

#### EXAMPLE 19

EXAMPLE 18 is repeated except that gabapentin of Example 4, having chloride ion of 7 ppm is used for formulating tablets. The resulting gabapentin tablet of  
25 each sample is initially measured to have 0.5 % by weight of lactam and approximately 7 ppm of chloride anion. The tablet is stored for one year at 25 °C and 60% atmospheric humidity and the increase in the lactam concentration is found not to exceed 0.2% by weight.

U.S. Patent Application

Filed: June 15, 2001

Inventors: Claude Singer, Gideon Pilarsky and Michael  
Pesachovitch

Attorney Docket No. 1662/52802

Examples 17-19 show that, contrary to Augart's disclosure, the presence of anion of a mineral acid in an amount greater than 20 ppm does not adversely affect the stability of gabapentin when stored for one year at 5 25 °C and 60% humidity (or higher). In addition, the Examples also show that gabapentin having pH in the range of 6.8 to 7.3, and preferably in the range of 7.0-7.2 is stable when stored for one year at 25 °C and 60% humidity. Further, the examples show that the 10 gabapentin formulations prepared in accordance with the invention showed equally stable result regardless of the type of adjuvant that were used.